

User's Manual and Instructions

QMTT Cell Viability Assay Kits (Z5030006, Z5030008)

Non-radioactive Colorimetric Assay for Cell Proliferation and Cytotoxicity

DESCRIPTION

The study of cell proliferation and cell viability requires the accurate quantification of the number of viable cells in a cell culture. Therefore, assays for calculating cell viability are necessary for optimizing cell culture conditions, evaluating cell growth factors and nutrients, discovering novel antibiotics and anti-cancer drugs, evaluating toxic effects of environmental pollutants and cell mediated toxicity and studying programmed cell death (apoptosis).

The QMTT assay kit provides a convenient, sensitive, quantitative and reliable assay for determining the number of viable cells in a given culture. This homogeneous colorimetric assay is based on the conversion of a tetrazolium salt MTT, a pale yellow substrate, to formazan, a purple dye. This cellular reduction reaction involves the pyridine nucleotide cofactors NADH/NADPH and is only catalyzed by living cells. The formazan product has a low aqueous solubility and is present as purple crystals. Dissolving the resulting formazan with a solubilization buffer permits the convenient quantification of product formation. The intensity of the product color, measured at 550 - 620 nm, is directly proportional to the number of living cells in the culture. Reagents in the kit have been carefully formulated and optimized for sensitivity, assay robustness and automation.

KEY FEATURES

Safe. Non-radioactive assay.

Sensitive and accurate. As low as 950 cells can be accurately quantified.

Convenient and high-throughput. "Mix-incubate-measure" type assay. No wash and reagent transfer steps are involved. Z' factors of 0.5 and above are observed. Can be readily automated with HTS liquid handling systems.

APPLICATIONS

Cell Proliferation: effects of cytokines, growth factor, nutrients.

Cytotoxicity and Apoptosis: evaluation of toxic compounds, anti-cancer antibodies, toxins, environmental pollutants etc.

Drug Discovery: high-throughput screen for toxic and anticancer drugs.

KIT CONTENTS

Reagent: 10mL

Solubilizer: 50mL

Control Reagent (Cat # Z5030008): 50 mg saponin (sold separately).

Storage conditions. Kit is shipped at room temperature. Store the Reagent at -20 °C. Store the Solubilizer at room temperature. Shelf life: 12 months.

Precautions: reagents are for research use only. Normal precautions for laboratory reagents should be exercised while using the reagents. Please refer to Material Safety Data Sheet for detailed information.

PROCEDURES

1. Plate and culture cells (80 μ L per well) in a clear bottom 96-well tissue culture plates. Typical culture medium contains DMEM, 10% fetal bovine serum, antibiotics (penicillin/streptomycin, gentamycin, etc), amino acids and other nutrients. Assays can be performed on either adherent cells or cells in suspension. The number of cells can vary from 1,000 to 80,000 per well. The volume can vary from 50 to 150 μ L, although 80 μ L is used in this example. In addition to the test samples, one must include control wells of culture medium containing no cells or cells treated with a toxic reagent such as 0.1% saponin.

2. Add test compounds and controls and incubate cells for the desired period of time (typically overnight). It is recommended that assays be run in duplicate or triplicate. A volume of 20 μ L in phosphate buffered saline (PBS) or culture medium is recommended for the test compounds and controls. The Control reagent can be conveniently reconstituted with 5 mL PBS (1% saponin).

3. Warm Reagent to room temperature. Add 15 μ L (per 80 μ L cell culture) of QMTT reagent per well and incubate for 4 hours at 37°C. The volume of the reagent should be adjusted depending on the volume of cell culture.

4. Add 100 μ L of the Solubilizer to each well. Mix gently on an orbital shaker for one hour at room temperature. The volume of the Solubilizer should be adjusted depending on the volume of cell culture. If precipitation occurs in the Solubilizer, place the bottle in a warm water bath or at 37°C and shake to dissolve precipitates.

5. Measure OD_{570nm} for each well on an absorbance plate reader. Maximum absorbance of the formazan dye lies between 560 and 590 nm. If desired, the OD measurement can be performed the following day. In this case, it is recommended to seal the plate to minimize evaporation.

DATA ANALYSIS

Determine the average of the blank controls and subtract this amount from all absorbance values. Plot the corrected absorbance values at 570nm against the concentration of the test compound. Determine the EC₅₀ value for cell proliferation and IC₅₀ value for cytotoxic compound by non-linear regression analysis using Prism or any other data analysis tools.

MATERIALS REQUIRED, BUT NOT PROVIDED

Pipetting devices, clear 96-well culture plates (e.g. VWR cat# 82050-760), and plate reader capable of either measuring absorbance between 560-590 nm.




Figure 1. HEK293 cells were plated in clear bottom 96-well plates. Cells were serially diluted from 55,000 cells to 25 cells in 100 μ L DMEM plus 10% FBS. 15 μ L of the QMTT reagent was added and cells were incubated for 4 hours at 37°C. Then 100 μ L of the Solubilizer was added and the plate was shaken for 1 hour at room temperature. A linear relationship was observed between OD_{570nm} and the cell number. The detection limit was estimated to be 950 cells from the blank control.




Figure 2. Cytotoxic effects of saponin in HEK293 cells. HEK293 cells were plated at 55,000 cells per 80 μ L well in clear bottom 96-well plate. Cells were treated with varying concentrations of saponin. After overnight incubation, 15 μ L of QMTT reagent was added and further incubated for 4 hours. 100 μ L of the Solubilizer was then added. The plate was shaken for 1 hour at room temperature. The IC₅₀ for saponin was 0.0026 wt%.

QMTT Cell Viability Assay Kits

Non-Radioactive Colorimetric Assay for Cell Proliferation and Cytotoxicity

LITERATURE

Cell proliferation

1. Luan X, Diekwisch TG (2002). CP27 affects viability, proliferation, attachment and gene expression in embryonic fibroblasts. *Cell Prolif.* 35: 207-19.

Cytotoxicity assays

2. Bezin C, Tomasi S, Lohezic-Le Devehat F, Boustie J (2003). Cytotoxic activity of some lichen extracts on murine and human cancer cell lines. *Phytomedicine.* 10(6-7):499-503.

3. Koh J, Kubota T, Koyama T, Migita T, Hashimoto M, Hosoda Y, Kitajima M (2000). Combined Antitumor Activity of 7-Hydroxystaurosporine (UCN-01) and Tamoxifen against Human Breast Carcinoma in vitro and in vivo. *Breast Cancer.* 10(3):260-7.

4. Labieniec M, Gabryelak T (2003). Effects of tannins on Chinese hamster cell line B14. *Mutat Res.* 539(1-2):127-35.

5. Camps J, About I (2003). Cytotoxicity testing of endodontic sealers: a new method. *J Endod.* 29(9):583-6.

6. Di Francesco AM, Mayalarp SP, Kim S, Butler J, Lee M (2003). Synthesis and biological evaluation of novel diaziridinylquinone-acridine conjugates. *Anticancer Drugs.* 14(8):601-15.

7. Moongkarndi P, Kaslungka S, Kosem N, Junnu S, Jongsomboonkusol S, Theptaranon Y, Neungton N (2003). Cytotoxicity and apoptosis of ovarian and breast cancer cell lines induced by OVS1 monoclonal antibody and paclitaxel. *Asian Pac J Allergy Immunol.* 21(1):31-41.

8. Yamashita T, Fujii M, Tomita T, Ishiguro R, Tashiro M, Tokumaru Y, Imanishi Y, Kanke M, Ogawa K, Kameyama K, Otani Y (2003). The inhibitory effect of matrix metalloproteinase inhibitor ONO-4817 on lymph node metastasis in tongue carcinoma. *Anticancer Res.* 23(3B):2297-302.

in vitro Chemosensitivity

9. Nakamori M, Iwahashi M, Nakamura M, Yamaue H(2003). Clinical benefit of chemosensitivity test for patients with regional lymph node-positive esophageal squamous cell carcinoma. *J Surg Oncol.* 84(1):10-6.

10. Ariyoshi Y, Shimahara M, Tanigawa N(2003). Study on chemosensitivity of oral squamous cell carcinomas by histoculture drug response assay. *Oral Oncol.* 39(7):701-7.

11. Yoshinare K, Kubota T, Watanabe M, Wada N, Nishibori H, Hasegawa H, Kitajima M, Takechi T, Fukushima M(2003). Gene expression in colorectal cancer and in vitro chemosensitivity to 5-fluorouracil: A study of 88 surgical specimens. *Cancer Sci.* 94(7):633-638.

12. Tallen G, Riabowol K, Wolff JE(2003). Expression of p33ING1 mRNA and chemosensitivity in brain tumor cells. *Anticancer Res.* 23(2B):1631-5.

Screening for cytotoxic compounds

13. Nishimura S, Murasugi T, Kubo T, Kaneko I, Meguro M, Marumoto S, Kogen H, Koyama K, Oda T, Nakagami Y(2003). RS-4252 inhibits amyloid beta-induced cytotoxicity in HeLa cells. *Pharmacol Toxicol.* 93(1):29-32.

14. Nakagami Y, Nishimura S, Murasugi T, Kubo T, Kaneko I, Meguro M, Marumoto S, Kogen H, Koyama K, Oda T(2002). A novel compound RS-0466 reverses beta-amyloid-induced cytotoxicity through the Akt signaling pathway in vitro. *Eur J Pharmacol.* 457(1):11-7.

15. Wu J, Zhang H, Wang J, Yang T, Xian J, Yang C, Zheng W, Chen H, Wang Q(2002). Screening for inhibitor of vascular endothelial growth factor from random peptide library. *Zhonghua Zhong Liu Za Zhi.* 24(6):540-3.

16. Ren DC, Du GH, Zhang JT(2003). High throughput screening for intercellular adhesion molecule-1 inhibitor. *Yao Xue Xue Bao.* 38(6):405-8.